FOR MEDICAL PROFESSIONALS

Differential Clinical Response in Oral Submucous Fibrosis (OSMF)

Clinical factors, differential diagnosis and structured reassessment during OSMF care

CLINICAL DISCUSSION • PATIENT-SPECIFIC RESPONSE • REASSESSMENT

Purpose. This resource outlines factors that may contribute to variable symptomatic and functional response in OSMF, highlights differential clinical possibilities, and provides a structured framework for reassessment. It is intended for professional education and clinical discussion.

01 Understanding differential clinical response

Oral Submucous Fibrosis (OSMF) is a chronic, progressive fibro-inflammatory disorder with variable biological behaviour. Clinical response can differ between patients because disease-related, behavioural, nutritional and patient-specific factors influence the course of rehabilitation.

A patient may report improvement in one domain—such as oral burning or eating comfort—while showing more limited change in another, such as inter-incisal mouth opening. Persistent symptoms do not, by themselves, establish that the underlying OSMF has worsened or that a care approach has failed. Equally, symptomatic improvement should not be taken as evidence that fibrosis or potentially malignant changes have resolved.

Clinical principle
Assess symptoms, oral function, mouth opening and mucosal findings separately. Compare current findings with a documented baseline and interpret them in the context of the individual patient.

02 Clinical domains to assess

Domain	What to document at review
Symptoms	Burning sensation, oral discomfort, soreness and tolerance of spicy food or a normal diet.
Function	Eating, speaking and other relevant oral functions; patient-reported quality of life.
Mouth opening	Current inter-incisal mouth opening in millimetres, measured consistently and compared with baseline.
Mucosal findings	Mucosal appearance, blanching, palpable fibrous bands, ulcers, plaques, erythematous or erosive areas, and any suspicious lesion.
Rehabilitation and adherence	Continuity of prescribed care and the frequency, quality and duration of mouth-opening exercises.

03 Factors that may influence response

Disease-related factors

- Stage and severity of OSMF.
- Duration of disease before treatment.
- Degree and extent of established fibrosis and reduced tissue elasticity.

- Baseline mouth opening.

Long-standing or advanced fibrosis may limit the extent and speed of functional improvement. In some patients, the practical objective may need to emphasise functional rehabilitation and quality of life rather than assuming complete reversal of established fibrosis.

Patient-related and behavioural factors

- Interrupted or inconsistent use of the prescribed care protocol (a “stop–start” pattern).
- Irregular, incorrectly performed or prematurely discontinued mouth-opening and jaw rehabilitation exercises.
- Continued or intermittent exposure to areca nut, gutkha, pan masala, tobacco or smoking.
- Nutritional status, oral hygiene, motivation and long-term follow-up.

Nutritional and systemic factors

The source material identifies iron, vitamin B12, folate and vitamin D deficiencies, diabetes and other systemic illnesses affecting tissue healing as factors for clinical consideration. Investigations should be selected according to the patient's history and clinical findings.

Coexisting oral conditions and local factors

Candidiasis, oral lichen planus, leukoplakia and other potentially malignant disorders, persistent ulceration, sharp teeth, ill-fitting dentures and chronic local trauma may independently contribute to burning, soreness or discomfort.

04 Differential clinical possibilities

Persistent burning, oral discomfort, mucosal changes or tissue stiffness should not automatically be attributed to OSMF alone. Coexisting disorders may produce overlapping features and may require independent diagnosis and management.

Condition / possibility	Features that may overlap or prompt consideration
Oral lichen planus (particularly erosive disease)	Burning, erythematous mucosa, erosive or ulcerative areas, soreness and discomfort with spicy food.
Chronic hyperplastic or atrophic candidiasis	Burning, soreness, altered taste and white plaques or other mucosal changes.
Leukoplakia and other potentially malignant disorders	White or otherwise altered mucosal lesions that require separate examination, documentation and appropriate evaluation.
Persistent ulceration / local trauma	Pain or burning associated with recurrent ulcers, sharp teeth, ill-fitting dentures or chronic mechanical irritation.
Scleroderma	May share features such as microstomia, reduced mouth opening, facial changes and dysphagia; systemic and skin findings and appropriate diagnostic criteria help distinguish it.
Amyloidosis	May present with hyalinized mucosa or stroma; systemic involvement and appropriate investigations, including special stains where indicated, may be relevant.
Iron deficiency anaemia	May be associated with mucosal pallor, burning, dysphagia and tongue changes; assess associated clinical features and haematological findings.
Generalized fibromatosis	May have histological features resembling fibrosis; clinical and histopathological assessment is required to distinguish it.

Do not overlook suspicious lesions
Persistent, indurated, unexplained, changing or otherwise suspicious lesions require timely clinical evaluation. Consider biopsy or specialist referral when clinically indicated. A patient with OSMF remains in need of appropriate surveillance even if symptoms improve.

05 Structured clinical reassessment

Before categorising a patient as a non-responder or concluding that care has been unsuccessful, consider a structured review of the following:

- Current inter-incisal mouth opening compared with baseline.
- Change in burning sensation, oral discomfort and tolerance of food.
- Eating, speech, oral function and quality-of-life concerns.
- Current OSMF stage, disease duration and extent of fibrosis.
- Adherence to the prescribed care protocol, including any interruptions.
- Frequency, technique and duration of prescribed jaw rehabilitation exercises.
- Current and past exposure to areca nut, gutkha, pan masala, tobacco and smoking; establish whether cessation is complete.
- Nutritional and systemic health assessment where clinically indicated.
- Examination for candidiasis, oral lichen planus, leukoplakia, traumatic lesions, persistent ulcers or other coexisting pathology.
- Identification of suspicious lesions requiring biopsy, further investigation or specialist referral.

Document change, not impression alone
Record baseline and follow-up measurements using a consistent method. Interpret objective findings alongside symptoms, function, mucosal examination and patient history.

06 Management considerations after reassessment

The next step should be guided by the clinical findings and the treating clinician's judgement. Depending on the case, considerations in the source documents include:

- Continuing or extending supportive rehabilitation where clinically appropriate.
- Reinforcing structured mouth-opening exercises and reviewing their technique.
- Addressing nutritional deficiencies or systemic conditions when identified.
- Managing coexisting oral diseases and eliminating local traumatic factors.
- Maintaining regular clinical review with serial mouth-opening measurements.
- Seeking specialist evaluation when the diagnosis is uncertain or malignant transformation is suspected.

MARCEL OMR Therapy™ is described in the supplied materials as a structured oral rehabilitation approach intended to support oral mucosal care and functional rehabilitation in OSMF. The initial care phase should not be treated as the sole determinant of outcome in every patient; any continuation or modification must be based on individual clinical assessment.

07 Case review information

For a case-specific discussion, the following information may help the clinical team understand the patient's course:

- Baseline and current inter-incisal mouth opening (mm).
- Clinical stage of OSMF and duration of symptoms.
- Habit history and current cessation status.
- Details of the care protocol followed, interruptions and adherence.
- Compliance with mouth-opening exercises.

- Clinical photographs and relevant examination findings.
- Relevant investigations, nutritional assessment or systemic history, where available.

Clinical information should be shared only with appropriate patient consent and through a suitable secure channel.

Clinical and safety note: This page is an educational resource for qualified healthcare professionals. It does not replace individual clinical examination, diagnosis, histopathological assessment where indicated, or professional judgement. Treatment decisions and follow-up should be based on the patient's findings. Suspicious lesions require timely evaluation.

JanSuta Pharmaceuticals
MARCEL OMR Therapy™ · For Medical Professionals